arXiv:1205.1121v1 [math.DS] 5 May 2012

GREEN FUNCTIONS AND WEIGHTS OF POLYNOMIAL SKEW PRODUCTS ON $\mathbb{C}^2$

KOHEI UENO

Abstract. We study the dynamics of polynomial skew products on $\mathbb{C}^2$. By using suitable weights, we prove the existence of several types of Green functions. Largely, continuity and plurisubharmonicity follow. Moreover, it relates to the dynamics of the rational extensions to weighted projective spaces.

1. Introduction

The Green function G_f of f is a powerful tool for the study of the dynamics of a polynomial map f from $\mathbb{C}^2$ to $\mathbb{C}^2$, which is defined as

$$G_f(z,w) = \lim_{n\to\infty} \frac{1}{\lambda^n} \log^+ |f^n(z,w)|,$$

where f^n denotes the n-th iterate of f, λ is the dynamical degree of f, and $|(z,w)| = \max\{|z|,|w|\}$. In one dimension, it is well known that the Green function G_p of a polynomial p is defined, continuous and subharmonic on $\mathbb{C}$ and that it coincides with the Green function of K_p with pole at infinity, where K_p is the set of points whose orbits under p are bounded. However, in two dimension, it is not known whether the limit G_f is defined on $\mathbb{C}^2$. One strategy to study the dynamics in two dimension is to assume that f has a good property that induces the existence of G_f. For example, it is well known that if f is regular, that is, if f extends to a holomorphic map on the projective space $\mathbb{P}^2$, then G_f is defined, continuous and plurisubharmonic on $\mathbb{C}^2$. Moreover, it coincides with the pluricomplex Green function of K_f, where K_f is the set of points whose orbits under f are bounded (see e.g. [1]). To relax this regularity condition, many authors have paid attention to another condition called algebraically stability (e.g. [11], [12], [7], [4], [5] and [6]). Recently, Favre and Jonsson [10] proved that any polynomial map has an extension that is algebraically stable in some weak sense, to a projective compactification of $\mathbb{C}^2$ with at worst quotient singularities. Their proof is based on the valuative techniques developed in [9], with

---

2010 *Mathematics Subject Classification.* 32H50 (primary), 30D05 (secondary).
Key words and phrases. Complex dynamics, skew product, Green function.

which they proved that if f is not conjugate to a skew product, then there is a plurisubharmonic function u, which is close to $\log^+ |(z,w)|$, such that $\lambda^{-n} u \circ f^n$ decreases to a plurisubharmonic function, which is not identically zero. Here $\circ$ denotes the composition. The distinctiveness of polynomial skew products was pointed out at numerous instances in their papers.

In this paper we study the dynamics of polynomial skew products on $\mathbb{C}^2$. A polynomial skew product is a polynomial map of the form $f(z,w) = (p(z), q(z,w))$. We assume that $\delta = \deg p \geq 2$ and $d = \deg_w q \geq 2$. Let $b(z)$ be the coefficient of w^d and γ its degree. Many authors have studied the dynamics of regular polynomial skew products (e.g. [13], [14], [15] and [3]); f is regular if and only if $\delta = d = \deg q$, which implies $\gamma = 0$. We have studied the dynamics of nondegenerate polynomial skew products in [17]; we say that f is nondegenerate if $\gamma = 0$. Besides giving examples of polynomial skew products whose Green functions are not defined on some curves, we introduced the weighted Green function G_f^α of f in [17],

$$G_f^\alpha(z,w) = \lim_{n \to \infty} \frac{1}{\lambda^n} \log^+ |f^n(z,w)|_\alpha,$$

where $|(z,w)|_\alpha = \max\{|z|^\alpha, |w|\}$, which is defined, continuous and plurisubharmonic on $\mathbb{C}^2$ for a suitable rational number $\alpha \geq 0$. Moreover, f extends to an algebraically stable map on a weighted projective space, whose dynamics relates to G_f^α. Although the dynamics becomes much more difficult without the nondegeneracy condition, the idea of imposing suitable weights is still effective.

The dynamics of f consists of the dynamics of p on the base space and the dynamics on fibers: $f^n(z,w) = (p^n(z), Q_z^n(w))$, where $Q_z^n = q_{p^{n-1}(z)} \circ \cdots \circ q_{p(z)} \circ q_z$ and $q_z(w) = q(z,w)$. As the Green function G_p of p, we define the fiberwise Green function of f as

$$G_z(w) = \lim_{n \to \infty} \frac{1}{d^n} \log^+ |Q_z^n(w)|.$$

Since G_p exists on $\mathbb{C}$, the existence of G_z implies the existence of G_f and G_f^α. Favre and Guedj proved the existence of G_z on $K_p \times \mathbb{C}$ in [8, Theorem 6.1], which is continuous and plurisubharmonic if $b^{-1}(0) \cap K_p = \emptyset$, and gave examples whose fiberwise Green functions are discontinuous over $J_p = \partial K_p$ in [8, Proposition 6.5]. In this paper, we investigate the existence, continuity and plurisubharmonicity of G_z on $A_p \times \mathbb{C}$, where A_p denotes the set of points whose orbits under p tend to infinity.

A summary of our results is as follows. We replace the definition of $|(z,w)|_\alpha$ by $\max\{|z|^{max\{\alpha,0\}}, |w|\}$ or $\max\{(|z|+1)^\alpha, |w|\}$. In the case

$\delta \neq d$, the weighted Green function G_f^α is defined on $\mathbb{C}^2$ and continuous and plurisubharmonic on $A_p \times \mathbb{C}$ for a suitable rational number α, which can be negative if $\delta < d$. If $\delta > d$, then α is nonnegative, $\limsup_{n\to\infty} \delta^{-n} \log^+ |Q_z^n| \leq \alpha G_p$ and so $G_f^\alpha = \alpha G_p$ on $\mathbb{C}^2$. Moreover, if $\alpha = \gamma/(\delta - d)$, then the limit

$$G_z^\alpha(w) = \lim_{n\to\infty} \frac{1}{d^n} \log^+ \left| \frac{Q_z^n(w)}{p^n(z)^\alpha} \right|$$

is defined, continuous and plurisubharmonic on $A_p \times \mathbb{C}$. If $\delta < d$, then G_z coincides with G_f and G_f^α on $\mathbb{C}^2$, and with G_z^α on $A_p \times \mathbb{C}$. Moreover, if $\delta \neq d$ and $\alpha = \gamma/(\delta - d)$, then we obtain certain uniform convergence to G_z^α on $A_p \times \mathbb{C}$ and the asymptotics of G_z^α near infinity. In the case $\delta = d$, the dynamics differs depending on whether f is nondegenerate. If $\gamma \neq 0$, then G_f^α is defined on $\mathbb{C}^2$ if we admit plus infinity for a suitable rational number α, which can be negative. Moreover, the limit $\lim_{n\to\infty} (n\gamma d^{n-1})^{-1} \log^+ |Q_z^n(w)|$ is defined and plurisubharmonic on $\mathbb{C}^2$. It coincides with G_p on a region in $A_p \times \mathbb{C}$ and with 0 on the complement of the region in $A_p \times \mathbb{C}$. Furthermore, the limit

$$G(z, w) = \lim_{n\to\infty} \frac{1}{d^n} \log \left| \frac{Q_z^n(w)}{p^n(z)^{n\frac{\gamma}{d}}} \right|$$

is defined and plurisubharmonic on $A_p \times \mathbb{C}$ if we admit minus infinity. It is continuous on the region in $A_p \times \mathbb{C}$.

Throughout the paper we make use of the extensions of f to rational maps on weighted projective spaces, which can be algebraically stable if and only if $\gamma = 0$ or $\delta > d$. Some results on the Green functions can be illustrated with these extensions. In particular, if $\delta > d$ and $\alpha = \gamma/(\delta - d)$ or if $\delta \leq d$, then $f \sim (z^\delta, z^\gamma w^d)$ on a region in the attracting basin of the indeterminacy point $[0 : 1 : 0]$, which induces the results on the Green functions. Here the notation $f \sim g$ means that the ratios of the first and second components of the maps tend to 1, respectively. Note that Guedj pointed out in [12, Example 3.2] that f extends to algebraically stable maps on Hirzebruch surfaces if $\delta \leq d$. Under an additional condition, he derived the existence of a weighted Green function of f in [12, Theorem 4.1], which is continuous and plurisubharmonic on $\mathbb{C}^2$.

This paper is organized as follows. First, we briefly recall the dynamics of polynomial skew products and give the definition of rational extensions on weighted projective spaces in Section 2. In Section 3 we recall the results for the nondegenerate case. The generalized results in the cases $\delta > d$, $\delta < d$ and $\delta = d$ are presented in Sections 4, 5 and 6, respectively. In these sections we offer two examples: monomial maps, and skew products that are semiconjugate to polynomial products. The weight α contributes also to the degree growth of f; in Section 7 we provide a result on the degree growth and a corollary on a weighted Green function that relates to the degree growth.

2. Preliminaries

Let $f(z,w) = (p(z), q(z,w))$ be a polynomial skew product such that $p(z) = az^\delta + O(z^{\delta-1})$ and $q(z,w) = b(z)w^d + O_z(w^{d-1})$, and let $\gamma = \deg b$. We assume that $\delta \geq 2$ and $d \geq 2$. Then we may assume that polynomials p and b are monic by taking an affine conjugate; $p(z) = z^\delta + O(z^{\delta-1})$ and $b(z) = z^\gamma + O(z^{\gamma-1})$. Let $\lambda = \max\{\delta, d\}$ and $\deg f$ be the algebraic degree of f, i.e. $\max\{\deg p, \deg q\}$. In general, $\deg f$ may be greater than λ. However, the dynamical degree of f, $\lim_{n\to\infty} \sqrt[n]{\deg(f^n)}$, is equal to λ.

Let us briefly recall the dynamics of polynomial skew products. Roughly speaking, the dynamics of f consists of the dynamics on the base space and the dynamics on the fibers. The first component p defines the dynamics on the base space $\mathbb{C}$. Note that f preserves the set of vertical lines in $\mathbb{C}^2$. For this reason, we often use the notation $q_z(w)$ instead of $q(z,w)$. The restriction of f^n to the vertical line $\{z\} \times \mathbb{C}$ can be viewed as the composition of n polynomials on $\mathbb{C}$, $q_{p^{n-1}(z)} \circ \cdots \circ q_{p(z)} \circ q_z$.

A useful tool in the study of the dynamics of p on the base space is the Green function G_p of p,

$$G_p(z) = \lim_{n\to\infty} \frac{1}{\delta^n} \log^+ |p^n(z)|.$$

It is well known that G_p is defined, continuous and subharmonic on $\mathbb{C}$. More precisely, G_p is harmonic and positive on A_p and zero on K_p, and $G_p(z) = \log|z| + o(1)$ as $z \to \infty$. Here $A_p = \{z : p^n(z) \to \infty\}$, $K_p = \{z : \{p^n(z)\}_{n\geq 1} \text{ bounded}\}$ and $A_p \sqcup K_p = \mathbb{C}$. By definition, $G_p(p(z)) = \delta G_p(z)$. In a similar fashion, we consider the fiberwise Green function of f,

$$G_z(w) = \lim_{n\to\infty} \frac{1}{d^n} \log^+ |Q_z^n(w)| \text{ or } G_z^\lambda(w) = \lim_{n\to\infty} \frac{1}{\lambda^n} \log^+ |Q_z^n(w)|,$$

where $Q_z^n = q_{p^{n-1}(z)} \circ \cdots \circ q_{p(z)} \circ q_z$. By definition, $G_{p(z)}(q_z(w)) = dG_z(w)$ if it exists. Since the limit G_p exists on $\mathbb{C}$, the existence of G_z or G_z^λ implies those of G_f and G_f^α. Since the existence of G_z on $K_p \times \mathbb{C}$ is proved in [8], the remaining problem lies in investigating the existence on $A_p \times \mathbb{C}$. We note that it is unclear even if f is regular. For the nondegenerate case, using an argument in the proof of [8, Theorem

6.1], the author showed the existence of G_z on an open subset of $A_p \times \mathbb{C}$ in [16, Lemma 2.3] with the assumption $\delta \leq d$, which was improved in [17]. In this paper we generalize these results for the case $\gamma \neq 0$.

It is useful to consider the dynamics of the rational extensions of f. Let r and s be any two positive integers. The weighted projective space $\mathbb{P}(r,s,1)$ is a quotient space of $\mathbb{C}^3 - \{O\}$,

$$\mathbb{P}(r,s,1) = \mathbb{C}^3 - \{O\} \ / \sim,$$

where $(z,w,t) \sim (c^r z, c^s w, ct)$ for any c in $\mathbb{C} - \{0\}$. Thus $\mathbb{P}(r,s,1) = \mathbb{C}^2 \sqcup L_\infty$, where L_∞ denotes the line at infinity $\{t = 0\}$. We denote a point in $\mathbb{P}(r,s,1)$ by weighted homogeneous coordinates $[z : w : t]$. It follows that f extends to a rational map $\tilde{f}$ on $\mathbb{P}(r,s,1)$,

$$\tilde{f}[z : w : t] = \left[p\left(\frac{z}{t^r}\right) t^{\lambda r} : q\left(\frac{z}{t^r}, \frac{w}{t^s}\right) t^{\lambda s} : t^\lambda \right].$$

We say that $\tilde{f}$ is algebraically stable if there is no integer n and no hypersurface V such that $\tilde{f}^n(V - I_{\tilde{f}^n}) \subset I_{\tilde{f}}$, where $I_{\tilde{f}}$ denotes the indeterminacy set of $\tilde{f}$. It is well known that if $\tilde{f}$ is algebraically stable on $\mathbb{P}^2$, then $\deg(f^n) = (\deg f)^n$. In the final section we present a similar claim on $\mathbb{P}(r,s,1)$. We define the Fatou set $F_{\tilde{f}}$ of $\tilde{f}$ as the maximal open set of $\mathbb{P}(r,s,1)$ where the family of iterates $\{\tilde{f}^n\}_{n \geq 0}$ is normal. The Julia set $J_{\tilde{f}}$ of $\tilde{f}$ is defined as the complement of the Fatou set of $\tilde{f}$.

3. Nondegenerate case

To recall the main results in [17], we assume that f is nondegenerate throughout this section. We defined the rational number α as

$$\min \left\{ l \in \mathbb{Q} \;\middle|\; \begin{array}{l} l\lambda \geq n_j + l m_j \text{ for any integers } n_j \text{ and } m_j \\ \text{s.t. } c_j z^{n_j} w^{m_j} \text{ is a term in } q \text{ for some } c_j \neq 0 \end{array} \right\}$$

if $\deg_z q > 0$ and as 0 if $\deg_z q = 0$. Since q has only finitely many terms, one can take the minimum. Indeed, α is equal to

$$\max \left\{ \frac{n_j}{\lambda - m_j} \;\middle|\; \begin{array}{l} c_j z^{n_j} w^{m_j} \text{ is a term in } q \\ \text{with } c_j \neq 0 \text{ and } m_j < \lambda \end{array} \right\}.$$

Clearly, $\alpha \geq 0$, and $\alpha = 0$ if and only if f is a polynomial product. By definition, $\alpha \leq \deg_z q$ and $\alpha < \deg q$. Moreover, $\lambda^n \leq \deg(f^n) \leq \max\{1, \alpha\}\lambda^n$ for any positive integer n.

We define the weight of a monomial $z^n w^m$ mainly as $n + \alpha m$. The weight of q is defined as the maximum of the weights of all terms in q. Then the weight of q is $\alpha\lambda$ and the weight of $Q_z^n(w)$ is $\alpha\lambda^n$. Let h be the weighted homogeneous part of q of highest weight $\alpha\lambda$. In the case $\delta \leq d$, it follows from definition that h contains w^d. On the other hand, if we replace α in the definition of the weight of a monomial by a positive number which is larger than α, then h coincides with w^d, and if we replace α by a positive number which is smaller than α, then h does not contain w^d.

We explain the importance of α in terms of the rational extensions of f, assuming that f is not a polynomial product. We saw that f extends to a rational map $\tilde{f}$ on $\mathbb{P}(r, s, 1)$ for any two positive integers r and s. Moreover, it extends to a weighted homogeneous polynomial on $\mathbb{C}^3$ if and only if $s/r \geq \alpha$. Similarly, $\tilde{f}$ is algebraically stable if and only if $s/r \geq \alpha$; if $s/r < \alpha$ then $\tilde{f}$ contracts $L_\infty - I_{\tilde{f}}$ to the indeterminacy point $p_\infty = [0 : 1 : 0]$. The map $\tilde{f}$ is holomorphic if and only if $\delta = d$ and $s/r \geq \alpha$. Note that the holomorphic extensions of polynomial maps to weighted projective spaces are also mentioned in [10, Section 5.3].

Let us list some more details of $\tilde{f}$ on $\mathbb{P}(r, s, 1)$ for $s/r \geq \alpha$, which deffer depending on the magnitude relation of δ and d. If $\delta > d$, then $I_{\tilde{f}} = \{p_\infty\}$. Moreover, if $s/r = \alpha$ then the dynamics on $L_\infty - \{p_\infty\}$ is induced by the polynomial h, and if $s/r > \alpha$ then $\tilde{f}$ contracts $L_\infty - \{p_\infty\}$ to the attracting fixed point $[1 : 0 : 0]$. On the other hand, p_∞ becomes an attracting fixed point if $\delta \leq d$. If $\delta < d$, then $\tilde{f}$ contracts $L_\infty - I_{\tilde{f}}$ to p_∞. If $\delta = d$, then the dynamics on L_∞ is determined by h. Here h is the weighted homogeneous part of q of highest weight sd/r, which coincides with w^d if $s/r > \alpha$.

Via α, we obtain the results on the Green functions of f. In the case $\delta > d$, the dynamics of $\tilde{f}$ described above implies the following upper estimate.

Theorem 3.1 ([17])**.** *Let $\gamma = 0$. If $\delta > d$, then $\tilde{G}_z^\lambda \leq \alpha G_p$ on $\mathbb{C}^2$, where $\tilde{G}_z^\lambda = \limsup_{n\to\infty} \delta^{-n} \log^+ |Q_z^n|$. In particular, if $\alpha \leq 1$ then $G_f = G_p$ on $\mathbb{C}^2$.*

Hence if $\delta > d$, then $G_f^\alpha = \alpha G_p$ on $\mathbb{C}^2$. In the case $\delta \leq d$, it follows from definition that $f \sim (z^\delta, w^d)$ on W_R and that $f(W_R) \subset W_R$ for large $R > 0$, where $W_R = \{|w| > R|z|^\alpha, |w| > R^{\alpha+1}\}$. This implies that G_z is defined, continuous and pluriharmonic on W_R. Although we use the same notation W_R in the sections below, the definition of W_R differs depending on whether f is nondegenerate. Let $A_f = \cup_{n\geq 0} f^{-n}(W_R)$.

Theorem 3.2 ([17])**.** *Let $\gamma = 0$. If $\delta \leq d$, then G_z is defined, continuous and pluriharmonic on A_f. Moreover, $G_z(w)$ tends to 0 if $\delta < d$ and to $\alpha G_p(z)$ if $\delta = d$ as (z, w) in A_f tends to ∂A_f.*

Hence if $\delta \leq d$, then G_f is defined, continuous and pluriharmonic on A_f. Note that A_f is the restriction of the attracting basin of p_∞ to $\mathbb{C}^2$. If $\delta < d$, then G_z is defined, continuous and plurisubharmonic on $\mathbb{C}^2$ and it coincides with both G_f and G_f^α. If $\delta = d$, then G_f^α is defined, continuous and plurisubharmonic on $\mathbb{C}^2$; roughly speaking, it is the maximum of αG_p and G_z. Moreover, G_f^α determines the Fatou and Julia sets of the holomorphic map $\tilde{f}$ on $\mathbb{P}(r,s,1)$, where $s/r \geq \alpha$.

Since two theorems above hold with suitable modifications for any $l \geq \alpha$, we arrive at the following corollary.

Corollary 3.3 ([17]). *Let $\gamma = 0$. For any $l \geq \alpha$, the limit G_f^l is defined, continuous and plurisubharmonic on $\mathbb{C}^2$.*

The α is optimal in the sense that in this theorem α can not be replaced by any smaller number. See [17, Remark 2, Examples 5.2 and 5.3] for details.

4. $\delta > d$

We first study the case $\delta > d$ toward a generalization of the results for nondegenerate polynomial skew products. Assume that f satisfies the condition $\delta > d$ throughout this section. Besides showing the same upper estimate of G_z as the nondegenerate case, we also analyze the new phenomena that does not appear in the nondegenerate case.

The definition of α is the same, which is positive unless f is a polynomial product. The map f extends to the rational map $\tilde{f}$ on $\mathbb{P}(r,s,1)$, which is algebraically stable if $s/r \geq \alpha$. The dynamics of $\tilde{f}$ implies the upper estimate of G_z on $\mathbb{C}^2$, which induces $G_f^\alpha = \alpha G_p$ on $\mathbb{C}^2$. Moreover, if $\alpha = \gamma/(\delta - d)$, then the dynamics of f is much more understandable; we show the existence of G_z^α on $A_p \times \mathbb{C}$. This function relates to the dynamics of $\tilde{f}$, and induces the existence of G_z and G_f on some region in $A_p \times \mathbb{C}$.

The organization of this section is as follows. In Section 4.1, we state the definition of α, whose importance is illustrated with the weighted homogeneous part of q or Q_z^n, and with the rational map $\tilde{f}$ on $\mathbb{P}(r,s,1)$. Then the upper estimate of G_z is described without a complete proof. In addition, we present two types of examples of polynomial skew products whose dynamics is rather understandable. The first type is monomial maps, whose Green functions are completely described. The second type is polynomial skew products that are semiconjugate to polynomial products, whose Green functions are well understood. With the assumption $\alpha = \gamma/(\delta - d)$, we prove the existence of G_z^α in Section 4.2

and the uniform convergence to G_z^α in Section 4.3, which implies the asymptotics of G_z^α near infinity.

4.1. **Weights.** The definition of α is the same as in the nondegenerate case:

$$\min \left\{ l \in \mathbb{Q} \;\middle|\; \begin{array}{l} l\delta \geq n_j + lm_j \text{ for any integers } n_j \text{ and } m_j \text{ s.t.} \\ z^{n_j} w^{m_j} \text{ is a term in } q \text{ with nonzero coefficient} \end{array} \right\},$$

which is equal to

$$\max \left\{ \frac{n_j}{\delta - m_j} \;\middle|\; \begin{array}{l} z^{n_j} w^{m_j} \text{ is a term in } q \\ \text{with nonzero coefficient} \end{array} \right\}.$$

Clearly, $\alpha \geq 0$, and $\alpha = 0$ if and only if f is a polynomial product. By definition, $\alpha \leq \deg_z q$ and $\alpha < \deg q$. Moreover, $\delta^n \leq \deg(f^n) \leq \max\{1, \alpha\}\delta^n$ for any positive integer n.

We define the weight of a monomial $z^n w^m$ mainly as $n + \alpha m$. As same as the nondegenerate case, the weight of q is $\alpha\delta$ and the weight of $Q_z^n(w)$ is $\alpha\delta^n$. Let h be the weighted homogeneous part of q of highest weight $\alpha\delta$, which contains $z^\gamma w^d$ if $\alpha = \gamma/(\delta - d)$ and does not if $\alpha > \gamma/(\delta - d)$. To begin with, let us consider the case where the rational number α is an integer. Put $w = cz^\alpha$, then $h(z, cz^\alpha) = h(1, c)z^{\alpha\delta}$. Fix c so that $q(z, cz^\alpha)$ can be regarded as a polynomial in z. By letting $h(c) = h(1, c)$, it follows from definition that $h(c)z^{\alpha\delta}$ is the homogeneous part of $q(z, cz^\alpha)$ of degree $\alpha\delta$. Moreover, $h^n(c)z^{\alpha\delta^n}$ is the homogeneous part of $Q_z^n(cz^\alpha)$ of degree $\alpha\delta^n$. Therefore, $z^{\alpha\delta^n} h^n(z^{-\alpha}w)$ is the weighted homogeneous part of $Q_z^n(w)$ of weight $\alpha\delta^n$. If α is not an integer, then z^α is not well defined and c is not uniquely determined by z and w. However, in that case, the polynomial h has some symmetries related to the denominator of α, and these notations are still helpful.

The dynamics of $\tilde{f}$ on $\mathbb{P}(r, s, 1)$ is also the same as the nondegenerate case. The details are as follows. If $s/r < \alpha$, then $\tilde{f}$ contracts $L_\infty - I_{\tilde{f}}$ to the indeterminacy point $p_\infty = [0 : 1 : 0]$; thus it is not algebraically stable. On the other hand, if $s/r \geq \alpha$, then $\tilde{f}$ is algebraically stable. If $s/r = \alpha$, then $\tilde{f}[z : w : t] = [z^\delta + tu(z, t) : h(z, w) + tv(z, w, t) : t^\delta]$, where u and v are polynomials. Since h is divisible by z, it follows that $I_{\tilde{f}} = \{p_\infty\}$. Since $\tilde{f}[z : w : 0] = [z^\delta : h(z, w) : 0]$, the dynamics of $\tilde{f}$ on $L_\infty - \{p_\infty\}$ is induced by the dynamics of $h(1, w)$. If $s/r > \alpha$, then $\tilde{f}[z : w : t] = [z^\delta + tu(z, t) : tv(z, w, t) : t^\delta]$. Hence $I_{\tilde{f}} = \{p_\infty\}$ and $\tilde{f}$ contracts $L_\infty - I_{\tilde{f}}$ to the attracting fixed point $[1 : 0 : 0]$. Because the fixed point $[1 : 0 : 0]$ is attracting in a very strong sense as above, we obtain the upper estimate $\tilde{G}_z^\lambda \leq lG_p$ on $A_p \times \mathbb{C}$ for any $l = s/r > \alpha$, where $\tilde{G}_z^\lambda = \limsup_{n \to \infty} \delta^{-n} \log^+ |Q_z^n|$. Therefore,

Proposition 4.1. *If $\delta > d$, then $\tilde{G}_z^\lambda \le \alpha G_p$ on $A_p \times \mathbb{C}$.*

One can also prove this proposition along the same line as the proof of [17, Theorem 3.2]. Hence if $\delta > d$, then $G_p \le \tilde{G}_f \le \max\{\alpha, 1\} G_p$ on $A_p \times \mathbb{C}$, where $\tilde{G}_f = \limsup_{n\to\infty} \delta^{-n} \log^+ |f^n|$. In particular, if $\alpha \le 1$, then $G_f = G_p$ on $\mathbb{C}^2$. In addition, the existence of G_z on $K_p \times \mathbb{C}$ implies that $G_f^\alpha = G_f = G_z^\lambda = 0$ on $K_p \times \mathbb{C}$, since $\lambda = \delta > d$. Consequently, the proposition above implies the following corollary.

Corollary 4.2. *If $\delta > d$, then $G_f^\alpha = \alpha G_p$ on $\mathbb{C}^2$.*

We end this subsection with two examples of polynomial skew products whose dynamics are well understood: monomial maps, and skew products that are semiconjugate to polynomial products.

Example 4.3 (monomial maps). *Let $f = (z^\delta, z^\gamma w^d)$, $\delta > d$ and $\gamma \neq 0$. Then $\alpha = \gamma/(\delta - d) > 0$ and $f^n = (z^{\delta^n}, z^{\gamma_n} w^{d^n})$, where*

$$\gamma_n = (\delta^{n-1} + \delta^{n-2} d + \cdots + d^{n-1})\gamma = \alpha \delta^n \left\{ 1 - \left(\frac{d}{\delta} \right)^n \right\}.$$

Hence G_z is ∞ on $\{|z| > 1, w \neq 0\}$, $\log^+ |w|$ on $\{|z| = 1\}$, and 0 on $\{|z| < 1\} \cup \{w = 0\}$. Since $G_p = \log^+ |z|$,

$$G_z^\lambda = \begin{cases} \alpha \log^+ |z| & \text{on } \{w \neq 0\} \\ 0 & \text{on } \{w = 0\}, \end{cases}$$

which is not continuous on $\{|z| \ge 1, w = 0\}$, and

$$G_f = \begin{cases} \max\{\alpha, 1\} \log^+ |z| & \text{on } \{w \neq 0\} \\ \log^+ |z| & \text{on } \{w = 0\}, \end{cases}$$

which is continuous on $\mathbb{C}^2$ if $\alpha \le 1$ and is not on $\{|z| \ge 1, w = 0\}$ if $\alpha > 1$. Therefore, $G_f^\alpha = \alpha \log^+ |z|$ on $\mathbb{C}^2$, which is continuous on $\mathbb{C}^2$. The limits G_z^λ, G_f and G_f^α are all plurisubharmonic on $\mathbb{C}^2$.

Example 4.4. *Let $f = (z^\delta, q(z, w))$ be a polynomial skew product, where $q = z^\gamma w^d + O_z(w^{d-1})$, $\gamma \neq 0$ and $\delta > d$, that is semiconjugate to $f_0 = (z^\delta, h(w))$ by $\pi = (z^r, z^s w)$ for some positive integers r and s; $f\pi = \pi f_0$. Note that $h(w) = q(1, w)$ and so the degree of h is d. The identity $q(z^r, z^s w) = z^{s\delta} q(1, w)$ implies that $\alpha = s/r > 0$ and $q(z, w) = z^{\alpha\delta} h(w/z^\alpha)$. Moreover, $f^n = (z^{\delta^n}, z^{\alpha\delta^n} h^n(w/z^\alpha))$. Hence $w_n/z_n^\alpha = h^n(w/z^\alpha)$ and so*

$$G_z^\alpha(w) = \lim_{n\to\infty} \frac{1}{d^n} \log^+ \left| \frac{w_n}{z_n^\alpha} \right| = G_h \left(\frac{w}{z^\alpha} \right) \text{ on } \mathbb{C}^2 - \{z = 0\},$$

where $(z_n, w_n) = f^n(z,w)$. *Define*

$$E_f = \bigcup_{|z|>1} \{z\} \times z^{\alpha} E_h \text{ and } E_h = \bigcap_{l \geq 0} \overline{\bigcup_{n \geq l} h^{-n}(0)}$$

as in [18]. *Then* $G_z^{\lambda} = \alpha \log^+ |z|$ *on* $\mathbb{C}^2 - E_f$, *which implies that* $G_f = \max\{\alpha, 1\} \log^+ |z|$ *on* $\mathbb{C}^2 - E_f$ *and that* $G_f^{\alpha} = \alpha \log^+ |z|$ *on* $\mathbb{C}^2$. *If* $0 \notin E_h$ *then the equalities above extend to* $\mathbb{C}^2$, *and if* $\alpha \leq 1$ *then* $G_f = \log^+ |z|$ *on* $\mathbb{C}^2$.

The claims in this example follow from the same line as in [18]. These maps can be characterized by the symmetries of the Julia sets, see [19, Theorems 5.2 and 5.5] for details.

4.2. **Existence of Green functions:** $\alpha = \gamma/(\delta - d)$**.** We saw that, for the special map f in Example 4.4, the limit G_z^{α} is defined, continuous and plurisubharmonic on $\mathbb{C}^2 - \{z = 0\}$. In this subsection, assuming that $\alpha = \gamma/(\delta - d)$, we derive the existence, continuity and plurisubharmonicity of G_z^{α} on $A_p \times \mathbb{C}$. We also assume that f is not a polynomial product for simplicity, which implies that $\alpha > 0$ and $\gamma \neq 0$.

Let $W_R = \{|z| > R, |w| > R|z|^{\alpha}\}$ for large $R > 0$. Note that the definition of W_R differs from the nondegenerate case. We often use the new variety $c = z^{-\alpha} w$. Although c depends on the choice of the branch of $z^{-\alpha}$, $|c|$ does not and $W_R = \{|z| > R, |c| > R\}$. The following important lemma follows from the definition of α.

Lemma 4.5. *If* $\delta > d$ *and* $\alpha = \gamma/(\delta - d)$, *then*

$$\left| \frac{q(z,w)}{p(z)^{\alpha}} \right| \sim \left| \frac{w}{z^{\alpha}} \right|^d \text{ on } W_R$$

and f *preserves* W_R*; that is,* $f(W_R) \subset W_R$.

Proof. We explain this claim by using the notation $|c| = |z^{-\alpha} w|$. Let $z^{n_j} w^{m_j}$ be a term of q with nonzero coefficient. If $z^{n_j} w^{m_j} \neq z^{\gamma} w^d$, then

$$\left| \frac{z^{n_j} w^{m_j}}{c^d z^{\alpha\delta}} \right| = \left| \frac{c^{m_j} z^{n_j + \alpha m_j}}{c^d z^{\alpha\delta}} \right| \to 0 \text{ as } z \text{ and } c \to \infty$$

since at least one of the inequalities $d > m_j$ and $\alpha\delta > n_j + \alpha m_j$ holds. Thus

$$\left| \frac{q(z,w)}{z^{\alpha\delta}} \right| = \left| \frac{q(z, cz^{\alpha})}{z^{\alpha\delta}} \right| = |c|^d \{1 + o(1)\} = \left| \frac{w}{z^{\alpha}} \right|^d \{1 + o(1)\}.$$

Hence there exist positive constants $r_1 < 1 < r_2$ such that

$$r_1 \left| \frac{w}{z^{\alpha}} \right|^d < \left| \frac{q(z,w)}{p(z)^{\alpha}} \right| < r_2 \left| \frac{w}{z^{\alpha}} \right|^d \tag{1}$$

on W_R for large R, since $p(z) \sim z^\delta$ as $z \to \infty$.

Let us show that f preserves W_R. It is clear that if R is large enough, then $|p(z)| > R$ since $p(z) \sim z^\delta$ as $z \to \infty$. Hence it is enough to show that $|q(z,w)| > R|p(z)|^\alpha$ on W_R for large R, which follows from inequality (1) since $d \geq 2$. $\square$

Remark 4.6. *We can also show the following as Lemmas 5.3 and 6.2 with a slight change of the proof: if $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then $q(z,w) \sim z^\gamma w^d$ on W_R for large $R > 0$.*

Let $\tilde{f}$ be the rational extension to $\mathbb{P}(r,s,1)$, where $s/r = \alpha$. Recall that the dynamics of $\tilde{f}$ restricted to $L_\infty - \{p_\infty\}$ is induced by the polynomial $h(1,w)$ of degree d. Hence p_∞ attracts most nearby points in $A_p \times \mathbb{C}$, and W_R is included in the attracting basin of p_∞. Therefore, the inclusion $f(W_R) \subset W_R$ is natural. Let $A_f = \cup_{n \geq 0} f^{-n}(W_R)$, which is the restriction of the attracting basin of p_∞ to $A_p \times \mathbb{C}$.

Theorem 4.7. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then the limit G_z^α is defined, continuous and pluriharmonic on A_f. Moreover, $G_z^\alpha = \log|z^{-\alpha}w| + o(1)$ on W_R, $G_z^\alpha \sim \log|w|$ as $w \to \infty$ for fixed z in A_p, and G_z^α tends to 0 as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$.*

Proof. First, we prove the uniform convergence of G_n to G_z^α on W_R, where $G_n(z,w) = d^{-n} \log|z_n^{-\alpha} w_n|$ and $(z_n, w_n) = f^n(z,w)$. It follows from inequality (1) in the proof of Lemma 4.5 that, for any (z,w) in W_R,

$$r_1 \left| \frac{w_n}{z_n^\alpha} \right|^d < \left| \frac{w_{n+1}}{z_{n+1}^\alpha} \right| < r_2 \left| \frac{w_n}{z_n^\alpha} \right|^d .$$

Hence, for any (z,w) in W_R and for any positive integer n,

$$|G_{n+1}(z,w) - G_n(z,w)| = \left| \frac{1}{d^{n+1}} \log \left| \frac{w_{n+1}}{z_{n+1}^\alpha} \right| \cdot \left| \frac{w_n}{z_n^\alpha} \right|^{-d} \right| < \frac{\log r}{d^{n+1}},$$

where $\log r = \max\{-\log r_1, \log r_2\}$. Therefore, G_n converges uniformly to G_z^α on W_R. Since G_n is continuous and pluriharmonic on W_R, the limit G_z^α is also continuous and pluriharmonic on W_R. By the inequality above, for any (z,w) in W_R,

$$\begin{aligned} \left| G_z^\alpha(w) - \log \left| \frac{w}{z^\alpha} \right| \right| &\leq \sum_{n=0}^{\infty} |G_{n+1}(z,w) - G_n(z,w)| \\ &< \sum_{n=0}^{\infty} \frac{\log r}{d^{n+1}} = \frac{\log r}{d-1} =: C_R. \end{aligned} \tag{2}$$

In particular, $G_z^\alpha \sim \log|w|$ as $w \to \infty$ for fixed z in W_R.

We can extend the domain of G_z^α from W_R to A_f. Indeed, for any (z,w) in A_f, there exists a positive integer n such that $f^n(z,w)$ belongs to W_R. Then we define $G_z^\alpha(w)$ as $d^{-n}G_{p^n(z)}^\alpha(Q_z^n(w))$. Clearly, G_z^α is continuous and pluriharmonic on A_f, and $G_z^\alpha \sim \log|w|$ as $w \to \infty$ for fixed z in A_p.

Next, we use inequality (2) to calculate the asymptotic value of $G_z^\alpha(w)$ as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$. Let $E = \{|w| = R|z|^\alpha, |z| > R\} \subset \partial W_R$. It then follows from inequality (2) that

$$\left| \, G_{p^n(z)}^\alpha(Q_z^n(w)) - \log R \, \right| \le C_R \text{ on } f^{-n}(E),$$

since $f^n(z,w)$ belongs to E for any (z,w) in $f^{-n}(E)$. Thus it follows from equation $G_{p^n(z)}^\alpha(Q_z^n(w)) = d^n G_z^\alpha(w)$ that

$$\left| \, G_z^\alpha(w) - d^{-n}\log R \, \right| \le d^{-n}C_R \text{ on } f^{-n}(E). \tag{3}$$

Since $f^{-n}(E)$ converges to $\partial A_f - J_p \times \mathbb{C}$ as n tends to infinity, $G_z^\alpha(w)$ converges to 0 as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$. □

We remark that the set $\partial A_f - J_p \times \mathbb{C}$ in Theorem 4.7 can be replaced by its closure. Let $B_f = A_p \times \mathbb{C} - A_f$. Since $G_z^\alpha = 0$ on B_f, we get the following corollary.

Corollary 4.8. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then the limit G_z^α is defined, continuous and plurisubharmonic on $A_p \times \mathbb{C}$. Moreover, it is pluriharmonic on A_f and intB_f.*

Here are some properties of A_f and B_f. First, A_f and B_f are invariant under f; that is, $f(A_f) \subset A_f = f^{-1}(A_f)$ and $f(B_f) \subset B_f = f^{-1}(B_f)$. Theorem 4.7 guarantees that the set $\{\{z\} \times \mathbb{C} : \deg_w q_z = 0\}$ of degenerate fibers does not intersect with A_f. Hence B_f is not empty; more precisely, $B_f \cap (\{z\} \times \mathbb{C}) \neq \emptyset$ for any z in A_p.

The existence of G_z^α implies the following two corollaries on the existence of other Green functions.

Corollary 4.9. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then $G_z = \infty$ on A_f.*

We note that there exist polynomial skew products such that $G_z = \infty$ on $A_p \times \mathbb{C}$. Indeed, let $f = (z^\delta, q(z,w))$ be a polynomial skew product that is semiconjugate to $(z^\delta, h(w))$ by $\pi = (z^r, z^s w)$. If $0 \in A_h$, then $G_z = \infty$ on $\{|z| > 1\}$.

Corollary 4.10. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then $G_z^\lambda = \alpha G_p$ and $G_f = \max\{\alpha, 1\} G_p$ on $\mathbb{C}^2 - B_f$.*

In particular, if $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then G_z^λ exists on A_f. We can insist on the optimality of α and A_f as in [17, Remark 2, Examples

5.2 and 5.3], using polynomial skew products that are semiconjugate to polynomial products.

We end this subsection with a description of the dynamics of $\tilde{f}$ on $\mathbb{P}(r,s,1)$, where $s/r = \alpha$. Let $U = \mathrm{int}\overline{A_p \times \mathbb{C}}$, where the interior and closure are taken in $\mathbb{P}(r,s,1)$. Then $U = (A_p \times \mathbb{C}) \cup (L_\infty - \{p_\infty\})$. Let $A_{\tilde{f}}$ be the restriction of the attracting basin of p_∞ to U. It is equal to the union of preimages $\tilde{f}^{-n}(\mathrm{int}\overline{W_R})$ and hence open. Moreover, $A_{\tilde{f}} = A_f \cup A_h$, where A_h denotes the set of points in $L_\infty - \{p_\infty\}$ whose orbits converge to p_∞. Since $A_{\tilde{f}}$ is included in the attracting basin of p_∞, it follows that $A_{\tilde{f}} \subset F_{\tilde{f}}$. Let $B_{\tilde{f}} = U - A_{\tilde{f}}$. Then $B_{\tilde{f}} = B_f \cup K_h$, where K_h denotes the set of points in $L_\infty - \{p_\infty\}$ whose orbits do not converge to p_∞. Since $\mathrm{int}B_{\tilde{f}} \cap \mathrm{int}\overline{\{|z| > R\}}$ is Kobayashi hyperbolic and preserved by $\tilde{f}$, it follows $\mathrm{int}B_{\tilde{f}} \subset F_{\tilde{f}}$ (see [17] for details).

Proposition 4.11. *Let $\delta > d$ and $\alpha = \gamma/(\delta - d)$. The restriction of $F_{\tilde{f}}$ to U consists of $A_{\tilde{f}}$ and int$B_{\tilde{f}}$. The restriction of $J_{\tilde{f}}$ to U is equal to the restriction of $\partial A_{\tilde{f}}$ to U, and to the restriction of $\partial B_{\tilde{f}}$ to U. Moreover, it coincides with the restriction of the closure of $\{(z,w) \in A_p \times \mathbb{C} : G_z^\alpha$ is not pluriharmonic$\}$ to U.*

The dynamics of $\tilde{f}$ on $B_{\tilde{f}}$ is as follows. Any point in $B_{\tilde{f}}$ is attracted to L_∞ under iteration. Eventually, the dynamics on L_∞, which is induced by h, should determine the dynamics of $\tilde{f}$ on $B_{\tilde{f}}$. It follows that $h(c) := h(1,c)$ can be written as $c^l H(c^r)$ for some integer $l \geq 0$ and some polynomial H, which is semiconjugate to $c^l H(c)^r$ by c^r. The restriction of $\tilde{f}$ to L_∞ is conjugate to $c^l H(c)^r$.

4.3. **Uniformly convergence and Asymptotics:** $\alpha = \gamma/(\delta - d)$.

Continuing with the assumption $\alpha = \gamma/(\delta - d)$, we show that the convergence to G_z^α is uniform on some region in $A_p \times \mathbb{C}$, which induces the asymptotic of G_z^α near infinity.

Proposition 4.12. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then the convergence to G_z^α is uniform on $V \times \mathbb{C}$, where $\overline{V} \subset A_p$.*

Proof. We show that, for any $\epsilon > 0$ and for any subset V such that $\overline{V} \subset A_p$, there exists N such that $|G_n - G_z^\alpha| < \epsilon$ on $V \times \mathbb{C}$ for any $n \geq N$, where $G_n(z,w) = d^{-n} \log^+ |z_n^{-\alpha} w_n|$ and $(z_n, w_n) = f^n(z,w)$. Let $U = \{z \in V, G_z^\alpha \geq \epsilon/3\}$. It is enough to show the uniform convergence of G_n to G_z^α on U. Indeed, if this holds, then there exists N such that $|G_n - G_z^\alpha| < \epsilon/3$ on U for any $n \geq N$. Hence $G_n < 2\epsilon/3$ on ∂U for any $n \geq N$. By Maximum Principle for subharmonic functions on vertical lines, $G_n < 2\epsilon/3$ and so $|G_n - G_z^\alpha| < \epsilon$ on U^c for any $n \geq N$, which completes the proof.

Now we show the uniform convergence on U. The equation $G_n \circ f = dG_{n+1}$ extends the uniform convergent region from W_R to $f^{-n}(W_R)$. Hence it is enough to show that $U \subset f^{-n}(W_R)$ for large n. From equation (3) in the proof of Theorem 4.7, it follows that $G_z^\alpha < \epsilon/3$ on $f^{-n}(E)$ for large n, which implies that $U \subset f^{-n}(W_R)$. □

We saw that, for fixed $c = z^{-\alpha}w$, the polynomial $h^n(c)z^{\alpha d^n}$ is the homogeneous part of $Q_z^n(cz^\alpha)$ of degree $\alpha\delta^n$. Although c is not well defined if α is not an integer, the polynomial h and the Green function G_h have some symmetries related to the denominator r of α in that case: $h(c)$ can be written as $c^l H(c^r)$, the Julia set J_h is preserved by the rotation ρc, where ρ is a r-th root of 1, and $G_h(c) = G_h(z^{-\alpha}w)$ is a well defined function in z and w. Hence we get the following asymptotics of G_z^α near infinity.

Lemma 4.13. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then $G_z^\alpha(cz^\alpha) = G_h(c) + o(1)$ as $z \to \infty$ for fixed c.*

Proof. We prove that for any $\epsilon > 0$ there exists $R > 0$ such that

$$|G_z^\alpha(cz^\alpha) - G_h(c)| < \epsilon \tag{4}$$

on $\{|z| > R\}$. Assume the rational number α is an integer. Since $Q_z^n(cz^\alpha) = h^n(c)z^{\alpha\delta^n} + o(z^{\alpha\delta^n})$, it follows that

$$\frac{1}{d^n}\log^+\left|\frac{Q_z^n(cz^\alpha)}{z^{\alpha\delta^n}}\right| = \frac{1}{d^n}\log^+|h^n(c)| + o(1).$$

Since $p(z) \sim z^\delta$ as $z \to \infty$,

$$\frac{1}{d^n}\log^+\left|\frac{Q_z^n(cz^\alpha)}{p^n(z)^\alpha}\right| = \frac{1}{d^n}\log^+|h^n(c)| + o(1). \tag{5}$$

By Proposition 4.12, there exist an integer N_1 and a number $R > 0$ such that $|G_z^\alpha(w) - G_n(z,w)| < \epsilon/3$ on $\{|z| > R\}$ for any $n \geq N_1$. Since the convergence of $G_h^n = d^{-n}\log^+|h^n|$ to G_h is uniform on $\mathbb{C}$, there exists N_2 such that $|G_h(c) - G_h^n(c)| < \epsilon/3$ for any $n \geq N_2$. From equation (5) it follows that if R is large enough, then $|G_N(z,w) - G_h^N(c)| < \epsilon/3$ on $\{|z| > R\}$, where $N = \max\{N_1, N_2\}$. Consequently, inequality (4) holds on $\{|z| > R\}$.

Even if α is not an integer, it follows that $|Q_z^n(cz^\alpha)| = |h^n(c)||z|^{\alpha\delta^n} + o(|z|^{\alpha\delta^n})$. A similar argument as above implies inequality (4). □

Proposition 4.14. *If $\delta > d$ and $\alpha = \gamma/(\delta - d)$, then $G_z^\alpha(w) = G_h(z^{-\alpha}w) + o(1)$ as $z \to \infty$.*

Proof. We prove that for any $\epsilon > 0$ there exists $R > 0$ such that

$$|G_z^\alpha(w) - G_h(z^{-\alpha}w)| < \epsilon \tag{6}$$

on $\{|z| > R\}$. Separating the region $\{|z| > R\}$ into two open sets whose union contains the region, we get this inequality.

First, we show that this inequality holds on W_R for large R, where $W_R = \{|z| > R, |w| > R|z|^\alpha\}$. Inequality (2) in the proof of Theorem 4.7 implies that $|G_z^\alpha(w) - \log|z^{-\alpha}w|| < \epsilon/2$ on W_R for large R. Since $|\log|z^{-\alpha}w| - G_h(z^{-\alpha}w)| < \epsilon/2$ on W_R for large R, there exists $R_1 > 0$ such that inequality (6) holds on W_{R_1}.

Next, let $V_R = \{|z| > R, |w| < 2R_1|z|^\alpha\}$. From a similar argument as the proof of Lemma 4.13, it follows that $G_z^\alpha(w) = G_h(z^{-\alpha}w) + o(1)$ on V_R, since the projection of V_R to $\mathbb{C}$ by the multi-valued function $z^{-\alpha}w$ is a relatively compact subset of $\mathbb{C}$. Hence there exists $R_2 \geq R_1$ such that inequality (6) holds on V_{R_2}. Since W_{R_2} and V_{R_2} cover the region $\{|z| > R_2\}$, inequality (6) holds on $\{|z| > R_2\}$. □

5. $\delta < d$

Next we study the case $\delta < d$, assuming that $\gamma \neq 0$. We generalize the definition of α, and prove the existence of G_z on $\mathbb{C}^2$ as the nondegenerate case, which is continuous and plurisubharmonic on $A_p \times \mathbb{C}$. Moreover, $G_z = G_f = G_f^\alpha$ on $\mathbb{C}^2$ and $G_z = G_z^\alpha$ on $A_p \times \mathbb{C}$. Unlike the nondegenerate case, α can be negative and $\tilde{f}$ is not algebraically stable unless f is nondegenerate. However, the indeterminacy point p_∞ is still attracting in some sense, and it follows that $f \sim (z^\delta, z^\gamma w^d)$ on a region in the attracting basin of p_∞, which induces the results above. To obtain uniform convergence to G_z, we have to consider unusual plurisubharmonic functions that converge to G_z. In particular, if $\alpha = \gamma/(\delta - d)$, then the convergence to G_z^α is uniform on a region in $A_p \times \mathbb{C}$, and we get the asymptotics of G_z^α near infinity.

This section is divided into three subsections: the definition and the properties of α, the existence of G_z and other Green functions, and the uniform convergence to G_z. We remark that many claims hold even if $\gamma = 0$.

5.1. **Weights.**

We generalize the definition of α as

$$\min\left\{ l \in \mathbb{Q} \;\middle|\; \begin{array}{l} \gamma + ld \geq l\delta \text{ and } \gamma + ld \geq n_j + lm_j \text{ for} \\ \text{any integers } n_j \text{ and } m_j \text{ s.t. } z^{n_j}w^{m_j} \text{ is} \\ \text{a term in } q \text{ with nonzero coefficient} \end{array} \right\},$$

which is equal to

$$\max\left\{ \frac{-\gamma}{d-\delta}, \frac{n_j-\gamma}{d-m_j} \;\middle|\; \begin{array}{l} z^{n_j}w^{m_j} \text{ is a term in } q \text{ with} \\ \text{nonzero coefficient s.t. } m_j < d \end{array} \right\}.$$

By definition, $-\gamma \leq \alpha \leq \deg_z q - \gamma$ and $\alpha < \deg q - \gamma$. Moreover, α induces inequalities for the degree growth of f, given in Section 7.

Let the weight of $z^n w^m$ be $n+\alpha m$. Then the weight of q is $\gamma+\alpha d$ and the weight of $Q_z^n(w)$ is $\gamma_n+\alpha d^n$, where $\gamma_n = (\delta^{n-1}+\delta^{n-2}d+\cdots+d^{n-1})\gamma$, which coincides with $\alpha\delta^n$ if $\alpha = \gamma/(\delta-d)$. Let h be the weighted homogeneous part of q of highest weight $\gamma+\alpha d$, which contains $z^\delta w^d$. If $\alpha = \gamma/(\delta-d)$, then $z^{\alpha\delta^n}h^n(z^{-\alpha}w)$ is the weighted homogeneous part of $Q_z^n(w)$ of weight $\alpha\delta^n$. If $\alpha > \gamma/(\delta-d)$, then $z^{\gamma_n}(z^{-\gamma}h(z,w))^{d^{n-1}}$ is the weighted homogeneous part of $Q_z^n(w)$ of weight $\gamma_n+\alpha d^n$.

The rational map $\tilde{f}$ on $\mathbb{P}(r,s,1)$ is not algebraically stable for any positive integers r and s; it contracts $L_\infty - I_{\tilde{f}}$ to the indeterminacy point p_∞. However, p_∞ attracts most points in $A_p \times \mathbb{C}$; more precisely, it attracts all points which converge to $(L_\infty - I_{\tilde{f}}) \cup \{p_\infty\}$.

As in Section 4.1, we end this subsection with two examples: monomial maps, and polynomial skew products that are semiconjugate to polynomial products.

Example 5.1 (monomial maps). *Let* $f = (z^\delta, z^\gamma w^d)$ *and* $\delta < d$. *Then* $\alpha = \gamma/(\delta-d) \leq 0$ *and* $f^n = (z^{\delta^n}, z^{\gamma_n}w^{d^n})$, *where*

$$\gamma_n = (\delta^{n-1}+\delta^{n-2}d+\cdots+d^{n-1})\gamma = -\alpha d^n \left\{1 - \left(\frac{\delta}{d}\right)^n\right\}$$

Hence $G_z = G_z^\alpha = G_f = G_f^\alpha = \log^+|z^{-\alpha}w|$, *which is continuous and plurisubharmonic on* $\mathbb{C}^2$.

Example 5.2. *Let* $f = (z^\delta, q(z,w))$ *be a polynomial skew product, where* $q = z^\gamma w^d + O_z(w^{d-1})$, $\gamma \neq 0$ *and* $\delta < d$, *that is semiconjugate to a polynomial product* $f_0 = (z^\delta, h(w))$ *by* $\pi = (z^r, z^{-s}w)$ *for some positive integers* r *and* s; $f\pi = \pi f_0$. *Note that* $h(w) = q(1,w)$, *which is divisible by* w. *It follows from the identity* $q(z^r, z^{-s}w) = z^{-s\delta}q(1,w)$ *that* $\alpha = -s/r < 0$ *and* $q(z,w) = z^{-\alpha\delta}h(z^{-\alpha}w)$. *Moreover,* $f^n = (z^{\delta^n}, z^{-\alpha\delta^n}h^n(z^{-\alpha}w))$. *Hence* $z_n^{-\alpha}w_n = h^n(z^{-\alpha}w)$ *and so*

$$G_z^\alpha(w) = \lim_{n\to\infty}\frac{1}{d^n}\log^+\left|z_n^{-\alpha}w_n\right| = G_h\left(z^{-\alpha}w\right) \text{ on } \mathbb{C}^2,$$

where $(z_n, w_n) = f^n(z,w)$. *Moreover,* $G_z = G_z^\alpha = G_f = G_f^\alpha$, *which is continuous and plurisubharmonic on* $\mathbb{C}^2$.

5.2. **Existence of Green functions.** We defined W_R as $\{|z| > R, |w| > R|z|^\alpha\}$ if $\gamma \neq 0$. The following important lemma follows from the definition of α.

Lemma 5.3. *If $\delta < d$, then $q(z,w) \sim z^\gamma w^d$ on W_R for large $R > 0$, and f preserves W_R; that is, $f(W_R) \subset W_R$.*

Proof. We explain this claim by using the notation $|c| = |z^{-\alpha}w|$. Let $z^{n_j}w^{m_j}$ be a term of q with nonzero coefficient. If $z^{n_j}w^{m_j} \neq z^\gamma w^d$, then

$$\left|\frac{z^{n_j}w^{m_j}}{z^\gamma w^d}\right| = \left|\frac{c^{m_j}z^{n_j+\alpha m_j}}{c^d z^{\gamma+\alpha d}}\right| \to 0 \text{ as } z \text{ and } c \to \infty$$

since at least one of the inequalities $d > m_j$ and $\gamma + \alpha d > n_j + \alpha m_j$ holds. Therefore, $q(z,w) \sim z^\gamma w^d$ on W_R. In other word,

$$r_1 < \left|\frac{q(z,w)}{z^\gamma w^d}\right| < r_2 \tag{7}$$

on W_R for some positive constants $r_1 < 1 < r_2$.

Let us show that f preserves W_R. It is enough to show that $|q(z,w)| > R|p(z)|^\alpha$ on W_R for large R. Since $p(z) \sim z^\delta$,

$$\left|\frac{q(z,w)}{p(z)^\alpha}\right| = \left|\frac{z^\delta}{p(z)}\right|^\alpha \cdot \left|\frac{z^{\gamma+\alpha d}}{z^{\alpha\delta}}\right| \cdot \left|\frac{q(z,w)}{z^{\gamma+\alpha d}}\right| \sim \left|\frac{z^{\gamma+\alpha d}}{z^{\alpha\delta}}\right| \cdot |c|^d.$$

This is larger than R if it is large enough since $\gamma + \alpha d \geq \alpha\delta$ and $d \geq 2$. ☐

This lemma implies that W_R is included in the attracting basin of p_∞. Let A_f be the union of preimages $f^{-n}(W_R)$.

Theorem 5.4. *If $\delta < d$, then the limit G_z is defined, continuous and pluriharmonic on A_f. Moreover, $G_z \sim \log|w|$ as $w \to \infty$ for fixed z in A_p, and G_z tends to 0 as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$.*

Proof. The proof is similar to that of Theorem 4.7. We first prove the locally uniform convergence of G_n to G_z on W_R, where $G_n = d^{-n}\log|Q_z^n|$. It follows from inequality (7) in the proof of Lemma 5.3 that, for any (z,w) in W_R,

$$r_1|(p^n(z))^\gamma| < \left|\frac{Q_z^{n+1}(w)}{Q_z^n(w)^d}\right| < r_2|(p^n(z))^\gamma|.$$

Hence, for any (z,w) in W_R and for any positive integer n,

$$|G_{n+1} - G_n| = \left|\frac{1}{d^{n+1}}\log\frac{|Q_z^{n+1}(w)|}{|Q_z^n(w)|^d}\right| < \frac{1}{d^{n+1}}\log\left(r|p^n(z)|^\gamma\right),$$

where $\log r = \max\{-\log r_1, \log r_2\}$. Therefore, G_n converges locally uniformly to G_z on W_R, which is continuous and pluriharmonic. By the inequality above, for any (z,w) in W_R,

$$|G_z(w) - \log|w|| < \sum_{n=0}^{\infty} \frac{\gamma}{d^{n+1}} \log|p^n(z)| + \frac{\log r}{d-1}.$$

Although this infinite sum converges since $\deg p = \delta < d$, we can restate this inequality to a simpler form. Since $p(z) \sim z^\delta$, there exists a constant $r_0 > 1$ such that $|p(z)| < r_0|z|^\delta$ and so $|p^n(z)| < (r_0|z|)^{\delta^n}$ if $|z| > R$. Hence

$$|G_z(w) - \log|w|| < \frac{\gamma}{d-\delta} \log|z| + C_R \tag{8}$$

for any (z,w) in W_R, where

$$C_R = \frac{\gamma}{d-\delta} \log r_0 + \frac{\log r}{d-1}.$$

In particular, $G_z \sim \log|w|$ as $w \to \infty$ for fixed z in W_R. We can extend the domain of G_z from W_R to A_f by the equation $G_z(w) = d^{-n} G_{p^n(z)}(Q_z^n(w))$.

Next, we calculate the asymptotic value of $G_z(w)$ as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$. By inequality (8),

$$\left| G_{p^n(z)}(Q_z^n(w)) - \log R|p^n(z)|^\alpha \right| \leq \frac{\gamma}{d-\delta} \log|p^n(z)| + C_R$$

for any (z,w) in $f^{-n}(E)$, where $E = \{|w| = R|z|^\alpha, |z| > R\} \subset \partial W_R$. Thus it follows from the equation $G_z(w) = d^{-n} G_{p^n(z)}(Q_z^n(w))$ that

$$\left| G_z(w) - \frac{1}{d^n} \log R|p^n(z)|^\alpha \right| \leq \frac{1}{d^n} \left(\frac{\gamma}{d-\delta} \log|p^n(z)| + C_R \right) \tag{9}$$

on $f^{-n}(E)$. Since $\deg p = \delta < d$ and since $f^{-n}(E)$ converges to $\partial A_f - J_p \times \mathbb{C}$ as n tends to infinity, $G_z(w)$ converges to 0 as (z,w) in A_f tends to $\partial A_f - J_p \times \mathbb{C}$. $\square$

Moreover, it follows that $G_z = \log|z^{\gamma/(d-\delta)} w| + o(1)$ on W_R; see Remark 5.8 in the next subsection. Since $G_z = 0$ on B_f, the theorem above implies the following.

Corollary 5.5. *If $\delta < d$, then G_z is defined on $\mathbb{C}^2$, which is continuous and plurisubharmonic on $A_p \times \mathbb{C}$. Moreover, $G_z = G_f = G_f^\alpha$ on $\mathbb{C}^2$, and $G_z = G_z^\alpha$ on $A_p \times \mathbb{C}$.*

We end this subsection with a description of the dynamics of $\tilde{f}$ on $\mathbb{P}(r,s,1)$ as in Section 4.2. By definition,

$$\begin{aligned}\tilde{f}[z:w:t] &= \left[t^{r(d-\delta)}\{z^\delta + tu(z,t)\} : \frac{h(z,w)+tv(z,w,t)}{t^{r\gamma}} : t^d\right] \\ &= \left[z^\delta + tu(z,t) : \frac{h(z,w)+tv(z,w,t)}{t^{r\gamma+sd-s\delta}} : t^\delta\right] \\ &= [t^{\frac{r}{s}A}\{z^\delta + tu(z,t)\} : h(z,w)+tv(z,w,t) : t^{\frac{1}{s}A+\delta}],\end{aligned}$$

where u and v are polynomials, h is the weighted homogeneous part of q of weight sd/r, and $A = r\gamma + sd - s\delta > 0$. The polynomial h contains $z^\gamma w^d$ if and only if $s/r \geq \alpha$, which coincides with $z^\gamma w^d$ if $s/r > \alpha$. Since $I_{\tilde{f}} = \{[z:w:0] : h(z,w) = 0\}$ and h is divisible by z, the point $p_\infty = [0:1:0]$ is always an indeterminacy point. Note that $\tilde{f}$ has other indeterminacy points than p_∞ if $s/r \geq \alpha$. In particular, $I_{\tilde{f}} = \{p_\infty, [1:0:0]\}$ if $s/r > \alpha$.

Recall that p_∞ attracts all points which converge to $(L_\infty - I_{\tilde{f}}) \cup \{p_\infty\}$. For the sets $A_{\tilde{f}}$ and $B_{\tilde{f}}$ as before, it follows that $A_{\tilde{f}} = A_f \cup (L_\infty - I_{\tilde{f}})$ and $B_{\tilde{f}} = B_f \cup (I_{\tilde{f}} - \{p_\infty\})$. Since $A_{\tilde{f}}$ is included in the attracting basin of p_∞ and since $B_{\tilde{f}}$ is the attracting basin of $I_{\tilde{f}} - \{p_\infty\}$, it follows that $A_{\tilde{f}} \cup \mathrm{int} B_{\tilde{f}} \subset F_{\tilde{f}}$. Hence the same claim as Proposition 4.11 holds.

5.3. **Uniformly convergence and Asymptotics.** It seems impossible to prove that the convergence of G_n to G_z is uniform, where $G_n = d^{-n} \log^+ |Q_z^n|$. To overcome this problem, we define

$$\tilde{G}_n(z,w) = G_n(z,w) + \sum_{j=n}^{\infty} \frac{\gamma}{d^{j+1}} \log^+ |p^j(z)|,$$

which converges to G_z on $\mathbb{C}^2$.

Proposition 5.6. *If $\delta < d$, then the convergence of $\tilde{G}_n$ to G_z is uniform on $V \times \mathbb{C}$, where $\overline{V} \subset A_p$.*

Proof. We first show that $\tilde{G}_n$ converges uniformly to G_z on W_R. By Lemma 5.3, there is a constant $r > 1$ such that, for any (z,w) in W_R,

$$\begin{aligned}\left|\tilde{G}_{n+1} - \tilde{G}_n\right| &= \left|G_{n+1} - G_n - \frac{\gamma}{d^{n+1}} \log |p^n(z)|\right| \\ &= \frac{1}{d^{n+1}} \log \left|\frac{Q_z^{n+1}(w)}{(p^n(z))^\gamma (Q_z^n(w))^d}\right| < \frac{\log r}{d^{n+1}}.\end{aligned}$$

Hence $\tilde{G}_n$ converges uniformly to G_z on W_R.

The left part of the proof is the same as the proof of Proposition 4.12. The equation $\tilde{G}_n \circ f = d\tilde{G}_{n+1}$ extends the uniform convergent region from W_R to $f^{-n}(W_R)$. It follows from equation (9) in the proof of Theorem 5.4 that $U = \{z \in V, G_z \geq \epsilon/3\} \subset f^{-n}(W_R)$ for large n, which induces the required uniform convergence. $\square$

This proposition holds even if we replace $\log^+ |p^j|$ in the definition of $\tilde{G}_n$ by $\log |p^j|$, since $|p^j| > 1$ on V if j is large enough. Moreover, in Proposition 5.6 we can replace $\tilde{G}_n$ by

$$\hat{G}_n = G_n + \sum_{j=n}^{\infty} \frac{\gamma}{d^{j+1}} \log |z^{\delta^j}| = G_n + \frac{\gamma}{d-\delta} \left(\frac{\delta}{d}\right)^n \log |z|$$

with a slight modification of the convergent region. However, the proof is not the same as before, because

$$\hat{G}_n \circ f - d\hat{G}_{n+1} = \frac{\gamma}{d-\delta} \left(\frac{\delta}{d}\right)^n \log \left|\frac{p(z)}{z^\delta}\right|,$$

which is not zero in general.

Proposition 5.7. *If $\delta < d$, then the convergence of $\hat{G}_n$ to G_z is uniform on $V \times \mathbb{C}$, where $\overline{V} \subset A_p \cap \{|z| > 1\}$.*

Proof. We first show that $\hat{G}_n$ converges uniformly to G_z on W_R. Since $p(z) \sim z^\delta$, there exists constants $0 < r_1 < 1 < r_2$ such that $r_1|z^\delta| < |p(z)| < r_2|z^\delta|$ and so $(r_1|z|)^{\delta^n} < |p^n(z)| < (r_2|z|)^{\delta^n}$ if $|z| > R$. Hence if $|z| > R$, then

$$\left|\log \left|\frac{p^n(z)}{z^{\delta^n}}\right|\right| < \log r_0^{\delta^n} = \delta^n \log r_0,$$

where $\log r_0 = \max\{-\log r_1, \log r_2\}$. By Lemma 5.3, there is a constant $r > 1$ such that, for any (z, w) in W_R,

$$\left|\log \left|\frac{Q_z^{n+1}(w)}{(p^n(z))^\gamma (Q_z^n(w))^d}\right|\right| < \log r.$$

With these inequalities, the equation

$$\left|\frac{Q_z^{n+1}(w)}{(z^{\delta^n})^\gamma (Q_z^n(w))^d}\right| = \left|\frac{p^n(z)}{z^{\delta^n}}\right|^\gamma \cdot \left|\frac{Q_z^{n+1}(w)}{(p^n(z))^\gamma (Q_z^n(w))^d}\right|$$

implies that, for any (z, w) in W_R,

$$\begin{aligned} \left|\hat{G}_{n+1} - \hat{G}_n\right| &= \left|G_{n+1} - G_n - \frac{\gamma}{d^{n+1}} \log |z^{\delta^n}|\right| \\ &= \left|\frac{1}{d^{n+1}} \log \left|\frac{Q_z^{n+1}(w)}{(z^{\delta^n})^\gamma (Q_z^n(w))^d}\right|\right| \qquad (10) \\ &< \frac{\gamma}{d} \left(\frac{\delta}{d}\right)^n \log r_0 + \frac{1}{d^{n+1}} \log r. \end{aligned}$$

Hence $\hat{G}_n$ converges uniformly to G_z on W_R.

Next, we show the uniform convergence on $f^{-N}(W_R) \cap (p_0^{-N}(U) \times \mathbb{C})$ for any N, where $p_0(z) = z^\delta$ and $U = \{|z| > R\}$. By definition, there exist constants $0 < m_N < 1 < M_N$ such that $m_N |z^{\delta^N}| < |p^N(z)| < M_N |z^{\delta^N}|$ on $U_N = p^{-N}(U) \cap p_0^{-N}(U)$. Since

$$\left| \frac{p^n(z)}{z^{\delta^n}} \right| = \left| \frac{p^{n-N}(p^N(z))}{p_0^{n-N}(p^N(z))} \cdot \frac{p_0^{n-N}(p^N(z))}{p_0^{n-N}(p_0^N(z))} \right|,$$

it follows that, for any z in U_N and for any $n \geq N$,

$$(r_1 m_N)^{\delta^{n-N}} < \left| \frac{p^n(z)}{z^{\delta^n}} \right| < (r_2 M_N)^{\delta^{n-N}}.$$

Applying a similar argument as above for $n \geq N$, we get the uniform convergence on the required region, which converges to $A_f \cap (\{|z| > 1\} \times \mathbb{C})$. The left part is the same as the proof of Proposition 4.12. □

Remark 5.8. *It follows from inequality* (10) *that*

$$\left| G_z(w) - \log |z^{\gamma/(d-\delta)} w| \right| < C_R \text{ on } W_R,$$

where C_R *converges to* 0 *as* R *tends to* ∞. *This inequality is better than inequality* (8) *in the proof of Theorem 5.4 and resembles inequality* (2) *in the proof of Theorem 4.7.*

Moreover, if $\alpha = \gamma/(\delta - d) < 0$, then we can show that the convergence to G_z^α is uniform. Let $\bar{G}_n = d^{-n} \log^+ |z_n^{-\alpha} w_n|$, where $(z_n, w_n) = f^n(z, w)$. If $\alpha = \gamma/(\delta - d) < 0$, then $\hat{G}_n = d^{-n} \log |z^{-\alpha\delta^n} w_n|$ and so

$$\bar{G}_n = \hat{G}_n + \frac{-\alpha}{d^n} \log \left| \frac{p^n(z)}{z^{\delta^n}} \right| \text{ on } W_R.$$

Hence we get the following convergence theorem.

Corollary 5.9. *If* $\delta < d$ *and* $\alpha = \gamma/(\delta - d) < 0$, *then the convergence of* $\bar{G}_n$ *to* G_z^α *is uniform on* $V \times \mathbb{C}$, *where* $\overline{V} \subset A_p$.

Proof. Since $\hat{G}_n$ converges uniformly to G_z^α on W_R, so does $\bar{G}_n$. The equation $\bar{G}_n \circ f = d\bar{G}_{n+1}$ extends the uniform convergent region from W_R to $f^{-n}(W_R)$. The left part is the same as the proof of Proposition 4.12. □

This uniform convergence induces the following asymptotics of G_z^α near infinity. The proofs are similar to those of Lemma 4.13 and Proposition 4.14.

Lemma 5.10. *If* $\delta < d$ *and* $\alpha = \gamma/(\delta - d)$, *then* $G_z^\alpha(cz^\alpha) = G_h(c) + o(1)$ *as* $z \to \infty$ *for fixed* c.

Proof. Because $Q_z^n(cz^\alpha) = z^{\alpha\delta^n} h^n(c)\{1 + o(1)\}$ as $z \to \infty$,

$$\hat{G}_n(z, cz^\alpha) = d^{-n} \log^+ |h^n(c)| + o(1).$$

Since $\hat{G}_n$ and $d^{-n} \log^+ |h^n|$ converge uniformly to G_z^α and G_h on suitable sets respectively, we get the required asymptotics. □

In this proof we can replace $\hat{G}_n$ by $\bar{G}_n$.

Proposition 5.11. *If $\delta < d$ and $\alpha = \gamma/(\delta - d)$, then $G_z^\alpha(w) = G_h(z^{-\alpha} w) + o(1)$ as $z \to \infty$.*

6. $\delta = d$

In this section we deal with the last case $\delta = d$. The results for the Green functions of f and the dynamics of $\tilde{f}$ are different depending on whether f is nondegenerate. However, it is common that p_∞ is attracting in some sense and that $f \sim (z^d, z^\gamma w^d)$ on a region in the attracting basin of p_∞.

In Section 6.1, we give the definition of α and an example of monomial maps. In Section 6.2, we prove the existence of three types of Green functions under the assumption $\gamma \neq 0$. First, we show that G_z is defined on A_f and G_f^α is defined on $\mathbb{C}^2$ if we admit plus infinity. Next, we show the existence of $\lim_{n\to\infty}(n\gamma d^{n-1})^{-1} \log^+ |Q_z^n(w)|$ on $\mathbb{C}^2$, which is continuous on $\mathbb{C}^2 - \partial A_f \cap \partial B_f$ and plurisubharmonic on $\mathbb{C}^2$. Finally, we show that the limit G is defined and plurisubharmonic on $A_p \times \mathbb{C}$ if we admit minus infinity. It is continuous and pluriharmonic on A_f.

6.1. **Weights.** We generalize the definition of α as

$$\inf \left\{ l \in \mathbb{Q} \;\middle|\; \begin{array}{l} \gamma + ld \geq n_j + lm_j \text{ for any integers } n_j \text{ and } m_j \text{ s.t.} \\ z^{n_j} w^{m_j} \text{ is a term in } q \text{ with nonzero coefficient} \end{array} \right\}.$$

This definition is similar to the previous case $\delta < d$, since the inequality $\gamma + ld \geq l\delta$ is trivial if $\delta = d$. If $q(z, w) \neq b(z)w^d$, then we can replace the infimum in the definition of α by the minimum, which is equal to

$$\max \left\{ \frac{n_j - \gamma}{d - m_j} \;\middle|\; \begin{array}{l} z^{n_j} w^{m_j} \text{ is a term in } q \text{ with} \\ \text{nonzero coefficient s.t. } m_j < d \end{array} \right\}.$$

For this case, $-\gamma \leq \alpha \leq \deg_z q - \gamma$ and $\alpha < \deg q - \gamma$. If $q(z, w) = b(z)w^d$, then $\alpha = -\infty$. Thus $\alpha = -\infty$ even if $q = w^d$, which differs with the definition we used for the nondegenerate case. See Section 7 for a claim regarding the degree growth of f.

Let the weight of $z^n w^m$ be $n + \alpha m$. Then the weight of q is $\gamma + \alpha d$ and the weight of $Q_z^n(w)$ is $n\gamma d^{n-1} + \alpha d^n$. Let h be the weighted homogeneous part of q of highest weight $\gamma + \alpha d$, which contains $z^{\delta} w^{d}$. If $\gamma = 0$, then $z^{\alpha \delta^n} h^n(z^{-\alpha} w)$ is the weighted homogeneous part of $Q_z^n(w)$ of weight αd^n. If $\gamma \neq 0$, then $z^{n\gamma d^{n-1}} (z^{-\gamma} h(z,w))^{d^{n-1}}$ is the weighted homogeneous part of $Q_z^n(w)$ of weight $n\gamma d^{n-1} + \alpha d^n$.

The dynamics of $\tilde{f}$ on $\mathbb{P}(r,s,1)$ is the same as in the case $\delta < d$ if $\gamma \neq 0$ for any positive integers r and s. Because it contracts $L_\infty - I_{\tilde{f}}$ to the indeterminacy point p_∞, the point p_∞ attracts most nearby points in $A_p \times \mathbb{C}$.

In this subsection we give only one example, i.e. monomial maps. If $\gamma = 0$, then there are many polynomial skew products that are semiconjugate to polynomial products; such maps are studied in [16, Theorem 3.7 and Proposition 3.9], [17, Examples 5.2 and 5.3] and [18]. However, we have no such maps if $\gamma \neq 0$.

Example 6.1 (monomial maps). *Let* $f = (z^d, z^\gamma w^d)$ *and* $\gamma \neq 0$. *Then* $\alpha = -\infty$ *and* $f^n = (z^{d^n}, z^{\gamma_n} w^{d^n})$, *where* $\gamma_n = n\gamma d^{n-1} + d^n$. *Hence*

$$G_f = G_z = \begin{cases} \infty & \text{on } \{|z| > 1, w \neq 0\} \\ \log^+ |w| & \text{on } \{|z| = 1\} \\ 0 & \text{on } \{|z| < 1\} \cup \{w = 0\}. \end{cases}$$

Moreover,

$$\lim_{n\to\infty} \frac{1}{\deg(f^n)} \log^+ |f^n(z,w)| = \lim_{n\to\infty} \frac{1}{n\gamma d^{n-1} + d^n} \log^+ |Q_z^n(w)|$$

$$= \begin{cases} \log^+ |z| & \text{on } \{w \neq 0\} \\ 0 & \text{on } \{w = 0\}, \end{cases}$$

which is plurisubharmonic on $\mathbb{C}^2$ *but not continuous on* $\{|z| \geq 1, w = 0\}$, *and* $G(z,w) = \log|w|$ *on* $\{z \neq 0\}$, *which is continuous and pluriharmonic on* $\{zw \neq 0\}$ *and plurisubharmonic on* $\{z \neq 0\}$.

6.2. **Existence of Green functions.** We defined W_R as $\{|z| > R, |w| > R|z|^\alpha\}$ if $\gamma \neq 0$. If $\alpha = -\infty$, then $W_R = \{|z| > R, |w| > 0\}$ since we may assume that $R > 1$. As same as the case $\delta < d$, we have the following lemma.

Lemma 6.2. *If* $\delta = d$, *then* $q(z,w) \sim z^\gamma w^d$ *on* W_R *for large* $R > 0$, *and* f *preserves* W_R; *that is,* $f(W_R) \subset W_R$.

Proof. If $q(z,w) \neq b(z) w^d$, then $\alpha > -\infty$ and the proof is the same as the case $\delta < d$, the proof of Lemma 5.3. If $q(z,w) = b(z)w^d$, then $\alpha = -\infty$ and this claim is trivial. Moreover, $q(z,w) \sim z^\gamma w^d$ on $\{|z| > R\} \times \mathbb{C}$. □

From now on, we deal with only the case $\gamma \neq 0$. The lemma above implies the following two theorems.

Theorem 6.3. *Let* $\delta = d$ *and* $\gamma \neq 0$. *If* $\alpha > 0$, *then* $G_z = \infty$ *on* A_f *and* $\tilde{G}_z \leq \alpha G_p$ *on* B_f.

Proof. By Lemma 6.2, there exists a positive constant $r < 1$ such that $|q(z,w)| > r|z^\gamma w^d|$ on W_R. Since $p(z) \sim z^d$, there exists a positive constant $r_0 < 1$ such that $|p(z)| > r_0|z|^d$ and so $|p^n(z)| > |r_0 z|^{d^n}$ if $|z| > R$. Using these inequalities inductively, we get

$$|Q_z^n(w)| > r^{1+d+\cdots+d^{n-1}} |(r_0 z)^{n\gamma d^{n-1}} w^{d^n}|$$

and so

$$\frac{1}{d^n} \log |Q_z^n(w)| > \log r + \frac{n\gamma}{d} \log |r_0 z| + \log |w|,$$

which tends to ∞ as $n \to \infty$. Therefore, $G_z = \infty$ on W_R, which extends to A_f.

Let (z,w) be a point in B_f. Then $f^n(z,w)$ never belong to W_R and so $|Q_z^n(w)| < R|p^n(z)|^\alpha$. Hence $\tilde{G}_z(w) \leq \alpha G_p(z)$ on B_f. □

The existence of G_z on B_f is still unclear. We exhibit three examples that relate to this problem.

Example 6.4. *For any positive integer* s, *let* $f = f_s = (z^2, z(w^2 - z^s) + z^{2s})$, *which is conjugate to* $f_1 = (z^2, zw^2)$ *by* $\pi = (z, w + z^s)$. *Then* $\alpha = s$ *and* $G_z = \alpha \log |z|$ *on* $B_f = \{|z| > 1, w = z^s\}$.

Example 6.5. *Let* $f = (z^r, z^\gamma(w^r - z^s) + z^s)$. *Then* $\alpha = s/r$ *and* $G_z = \alpha \log |z|$ *on* $\{|z| > 1, w^r = z^s\} \subset B_f$.

Example 6.6. *Let* $f = (z^2, zw^2 + z^2 w)$. *Then* $\alpha = 1$ *and* $G_z = 0$ *on* $\{w = 0\} \subset B_f$.

Let us return to the statement on the Green function in the remaining case $\alpha \leq 0$.

Theorem 6.7. *Let* $\delta = d$ *and* $\gamma \neq 0$. *If* $\alpha \leq 0$, *then* G_z *is* ∞ *on* A_f *and* 0 *on* B_f.

Proof. If $q(z,w) \neq b(z)w^d$, then $\alpha > -\infty$. The proof of the claim $G_z = \infty$ on A_f is the same as the proof of Theorem 6.3. The claim $G_z = 0$ on B_f follows from the definition of B_f and the assumption $\alpha \leq 0$. If $q(z,w) = b(z)w^d$, then $\alpha = -\infty$ and this claim follows from the direct calculation; see the proof of Proposition 6.8 below for detail. □

More precisely, B_f consists of infinitely many lines if $q = b(z)w^d$.

Proposition 6.8. *Let* $\delta = d$ *and* $\gamma \neq 0$*. If* $q(z,w) = b(z)w^d$*, then* G_z *is* ∞ *on* A_f *and* 0 *on* B_f*. Moreover,* B_f *coincides with the union of the preimages of* $\{z \in A_p, w = 0\}$ *under* f*, which is equal to*

$$\bigcup_{n \geq 0} p^{-n}(b^{-1}(0) \cap A_p) \times \mathbb{C} \cup \{z \in A_p, w = 0\}.$$

Proof. Let $f(z,w) = (p(z), b(z)w^d)$. Then $f^n(z,w) = (p^n(z), B_n(z)w^{d^n})$, where $B_n(z) = b(p^{n-1}(z)) \cdots b(p(z))^{d^{n-2}} b(z)^{d^{n-1}}$. Hence

$$\frac{1}{d^n} \log |Q_z^n(w)| = \log |w| + \sum_{j=0}^{n-1} \frac{1}{d^{j+1}} \log |b(p^j(z))|.$$

This finite sum tends to ∞ as $n \to \infty$ unless $b(p^j(z)) = 0$ for some $j \geq 0$, since $\deg p = d$. $\square$

Combining two theorems above, we get the following corollary.

Corollary 6.9. *If* $\delta = d$ *and* $\gamma \neq 0$*, then*

$$G_f^\alpha(z,w) = \begin{cases} \infty & \text{on } A_f \\ \max\{\alpha, 0\} G_p(z) & \text{on } B_f. \end{cases}$$

The dynamics of $\tilde{f}$ is similar to that mentioned in the previous section. The set $A_{\tilde{f}}$ is included in the attracting basin of p_∞, and $B_{\tilde{f}}$ is the attracting basin of $I_{\tilde{f}} - \{p_\infty\}$. Hence the same claim as Proposition 4.11 holds except the description of $J_{\tilde{f}}$ in terms of a Green function, for which we use the Green function in Corollary 6.11 below instead of G_z^α.

Now we show the existence of other Green functions that are locally bounded on $\mathbb{C}^2$.

Theorem 6.10. *If* $\delta = d$ *and* $\gamma \neq 0$*, then*

$$\lim_{n \to \infty} \frac{1}{n\gamma d^{n-1}} \log |Q_z^n(w)| = G_p(z) \ \text{on } A_f.$$

Proof. By Lemma 6.2, there exist constants $r_1 < 1 < r_2$ such that $r_1 |z^\gamma w^d| < |q(z,w)| < r_2 |z^\gamma w^d|$ on W_R. By using $|q(z,w)| < r_2 |z^\gamma w^d|$ inductively, we get the upper estimate

$$|Q_z^n(w)| < r_2^{1+d+\cdots+d^{n-1}} |p^{n-1}(z)|^\gamma |p^{n-2}(z)|^{\gamma d} \cdots |z|^{\gamma d^{n-1}} |w|^{d^n}$$

and so

$$\frac{1}{n\gamma d^{n-1}} \log |Q_z^n(w)| < \frac{d}{n\gamma} (\log r_2 + \log |w|)$$
$$+ \frac{1}{n} \left\{ \log |z| + \frac{1}{d} \log |p(z)| + \cdots + \frac{1}{d^{n-1}} \log |p^{n-1}(z)| \right\}.$$

Since $d^{-n}\log|p^n|$ converges to G_p on A_p, the right hand side converges to G_p as $n\to\infty$. Thus the inequality

$$\limsup_{n\to\infty}\frac{1}{n\gamma d^{n-1}}\log|Q_z^n(w)|\le G_p(z)$$

holds on W_R. By the same argument in terms of $r_1|z^\gamma w^d|<|q(z,w)|$, we get the inverse inequality. Therefore, we get the required equation on W_R.

A similar argument as above induces the required equation on A_f, because if $f^N(z,w)$ belongs to W_R then $Q_z^n(w)=Q_{p^N(z)}^{n-N}(Q_z^N(w))$ approximates to $p^{n-1}(z)^\gamma p^{n-2}(z)^{\gamma d}\cdots p^N(z)^{\gamma d^{n-N-1}}Q_z^N(w)^{d^{n-N}}$ for $n\ge N$. □

It follows from this theorem that $\lim_{n\to\infty}(n\gamma d^{n-1})^{-1}\log^+|Q_z^n|=0$ on B_f. Therefore,

Corollary 6.11. *If $\delta=d$ and $\gamma\ne 0$, then*

$$\lim_{n\to\infty}\frac{1}{n\gamma d^{n-1}}\log^+|f^n(z,w)|=\lim_{n\to\infty}\frac{1}{n\gamma d^{n-1}}\log^+|Q_z^n(w)|$$

$$=\begin{cases}G_p(z) & \text{on } \mathbb{C}^2-B_f\\ 0 & \text{on } B_f.\end{cases}$$

Finally, we prove the existence of G using Lemma 6.2. By definition, $G(f^n(z,w))=d^nG(z,w)+n\gamma d^{n-1}G_p(z)$ if it exists.

Theorem 6.12. *The limit G is defined, continuous and pluriharmonic on A_f. Moreover, $G=\log|w|+o(1)$ on W_R, $G\sim\log|w|$ as $w\to\infty$ for fixed z in A_p, and G tends to $-\infty$ as (z,w) in A_f tends to any point in $\partial A_f-J_p\times\mathbb{C}$.*

Proof. The proof is similar to those of Theorems 4.7 and 5.4. We first show the uniform convergence to G on W_R. Let $G_n=d^{-n}\log|z_n^{-n\gamma/d}w_n|$, where $(z_n,w_n)=f^n(z,w)$. By Lemma 6.2, there are constants $r>1$ and $r_0>1$ such that $|w_{n+1}|<r|z_n^\gamma w_n^d|$ and $|z_n^d|<r_0|z_{n+1}|$ on W_R. Hence

$$G_{n+1}-G_n=\frac{1}{d^{n+1}}\log\left|\frac{w_{n+1}}{w_n^d}\cdot\frac{z_n^{n\gamma}}{z_{n+1}^{(n+1)\frac{\gamma}{d}}}\right|<\frac{1}{d^{n+1}}\log\left|r\left(\frac{z_n^d}{z_{n+1}}\right)^{(n+1)\frac{\gamma}{d}}\right|$$

$$<\frac{1}{d^{n+1}}\log\left|r{r_0}^{(n+1)\frac{\gamma}{d}}\right|=\frac{1}{d^{n+1}}\log r+\frac{n+1}{d^{n+1}}\cdot\frac{\gamma}{d}\log r_0.$$

Therefore, G_n converges uniformly to G on W_R, which is continuous and pluriharmonic. By the inequality above,

$$|G(z,w)-\log|w||<C_R \text{ on } W_R, \tag{11}$$

where the constant $C_R > 0$ converges to 0 as $R \to \infty$. The equation $G = d^{-n} G \circ f^n - n\gamma d^{-1} G_p$ extends the domain of G from W_R to A_f.

Next, we show the last statement. By inequality (11),

$$|G(f^n(z,w)) - \log R|p^n(z)|^\alpha| < C_R \text{ on } f^{-n}(E),$$

where $E = \{|w| = R|z|^\alpha, |z| > R\} \subset \partial W_R$. Thus, by the identity $G \circ f^n = d^n G + n\gamma d^{n-1} G_p$,

$$\left| G(z,w) + n\frac{\gamma}{d} G_p(z) - \frac{\log R|p^n(z)|^\alpha}{d^n} \right| < \frac{C_R}{d^n} \text{ on } f^{-n}(E).$$

Therefore, the values of G on $f^{-n}(E)$ tend to $-\infty$ as $n \to \infty$. $\square$

This theorem guarantees that $B_f \cap (\{z\} \times \mathbb{C}) \neq \emptyset$ for any z in A_p. Since $G = -\infty$ on B_f, it follows that G is defined and plurisubharmonic on $A_p \times \mathbb{C}$, and $A_f = \{(z,w) \in A_p \times \mathbb{C} : G > -\infty\}$. The identity $G \circ f^n = d^n G + n\gamma d^{n-1} G_p$ induces that $G(z,w)$ converges to 0 as (z,w) in A_f tends to the intersection of the closure of $\partial A_f - J_p \times \mathbb{C}$ and $J_p \times \mathbb{C}$. Hence the description of the last statement is different from those in the case $\delta \neq d$.

7. Degree growth

In this section we give a remark on the relationship between algebraically stability and weight growth, the list of inequalities of degree growth in terms of α, and a corollary on the existence of a weighted Green function that is normalized by $\deg(f^n)$.

7.1. **Algebraically stability and Weight growth.** Assume that f is not a polynomial product. We saw that if $\gamma \neq 0$ or $\delta > d$, then α is a positive rational number, f extends to an algebraically stable map on $\mathbb{P}(r,s,1)$, where $s/r = \alpha$, and the weight of Q_z^n is $\alpha \lambda^n$. Hence the weight of f^n is equal to λ^n if we define it as the maximum of the degree of p^n and the weight of Q_z^n times α^{-1}; namely, $\text{weight}(f^n) = (\text{weight} f)^n$. This is an analogue of the well known fact: if $\tilde{f}$ is algebraically stable on $\mathbb{P}^2$, then $\deg(f^n) = (\deg f)^n$.

7.2. **Degree growth.** Next, we give the list of inequalities on the degree growth of f, which follows from the definition of α.

(1) If $\delta > d$, then $\delta^n \leq \deg(f^n) \leq \max\{\alpha, 1\}\delta^n$.

(2) If $\delta < d$ and $\alpha > 0$, then

$$\left[1 + \frac{\gamma}{d-\delta} \left\{ 1 - \left(\frac{\delta}{d} \right)^n \right\} \right] d^n \leq \deg(f^n)$$

$$\leq \left[\max\{\alpha, 1\} + \max\left\{\frac{1}{\alpha}, 1\right\} \frac{\gamma}{d-\delta}\left\{1 - \left(\frac{\delta}{d}\right)^n\right\}\right] d^n.$$

If $\delta < d$ and $\alpha \leq 0$, then

$$\deg(f^n) = \left[1 + \frac{\gamma}{d-\delta}\left\{1 - \left(\frac{\delta}{d}\right)^n\right\}\right] d^n.$$

(3) If $\delta = d$ and $\alpha > 0$, then

$$\left(\frac{\gamma}{d}n + 1\right) d^n \leq \deg(f^n) \leq \left[\max\left\{\frac{1}{\alpha}, 1\right\} \frac{\gamma}{d}n + \max\{\alpha, 1\}\right] d^n.$$

If $\delta = d$ and $\alpha \leq 0$, then

$$\deg(f^n) = \left(\frac{\gamma}{d}n + 1\right) d^n.$$

Note that it follows from the equalities above that, if $\gamma = 0$, then $\lambda^n \leq \deg(f^n) \leq \max\{\alpha, 1\}\lambda^n$, which was already stated in [17]. In general, it is proved in [9] that, for any polynomial map F which is not conjugate to a skew product such that $\delta = d$ and $\gamma \neq 0$, if the dynamical degree λ is larger than 1, then there is $D \geq 1$ such that $\lambda^n \leq \deg(F^n) \leq D\lambda^n$.

7.3. **Another Green function.**

In the final subsection we consider the existence of the limit

$$\lim_{n\to\infty} \frac{1}{\deg(f^n)} \log^+ |f^n(z,w)|_\alpha.$$

If $\delta \neq d$, then the topological degree δd is smaller than λ^2. In this case, the existence of the limit follows from Corollaries 4.2 and 5.5 and the main result in [2], which can be restated as follows: for a polynomial map F whose topological degree is smaller than the square λ^2 of the dynamical degree, the ratio of $\deg(F^n)$ and λ^n converges to some positive number. For the case $\delta = d$, the existence follows if $\alpha \leq 0$ because of Corollary 6.11 and the result above on the degree growth. Consequently,

Corollary 7.1. *If $\delta \neq d$ or $\alpha \leq 0$, then the limit of $(\deg(f^n))^{-1} \log^+ |f^n(z,w)|_\alpha$ is defined on $\mathbb{C}^2$. Moreover, it is continuous on $\mathbb{C}^2$ if $\delta > d$, on $A_p \times \mathbb{C}$ if $\delta < d$, and on $\mathbb{C}^2 - B_f$ if $\delta = d$ and $\alpha \leq 0$.*

REFERENCES

[1] E. Bedford and M. Jonsson. Dynamics of regular polynomial endomorphisms of $\mathbb{C}^k$. *American J. Math.* **122** (2000), 153-212.

[2] S. Bouckson, C. Favre and M. Jonsson. Degree growth of meromorphic surface maps. *Duke. Math. J.* **141** (2008), 519-538.

[3] L. DeMarco and S. L. Hruska. Axiom A polynomial skew products of $\mathbb{C}^2$ and their postcritical sets. *Ergodic Theory Dynam. Syst.* **28** (2008), 1729-48.

[4] J. Diller, R. Dujardin and V. Guedj. Dynamics of meromorphic maps with small topological degree I: from cohomology to currents. *Indiana Univ. Math. J.* **59** (2010), 521-561.

[5] J. Diller, R. Dujardin and V. Guedj. Dynamics of meromorphic maps with small topological degree II: Energy and invariant measure. *Comment. Math. Helv.* **86** (2011), 277-316.

[6] J. Diller, R. Dujardin and V. Guedj. Dynamics of meromorphic maps with small topological degree III: geometric currents and ergodic theory. *Ann. Sci. Ec. Norm. Super.* **43** (2010), 235-278.

[7] J. Diller and V. Guedj. Regularity of dynamical Green's functions. *Transactions of the Amer. Math. Soc.* **361** (2009), 4783-4805.

[8] C. Favre and V. Guedj. Dynamique des applications rationnelles des espaces multiprojectifs. *Indiana Univ. Math. J.* **50** (2001), 881-934.

[9] C. Favre and M. Jonsson. Eigenvaluations. *Ann. Sci. École Norm. Sup.* **40** (2007), 309-349.

[10] C. Favre and M. Jonsson. Dynamical compactifications of $\mathbb{C}^2$. *Ann. of Math.* **173** (2011), 211-248.

[11] J-E. Fornaess and N. Sibony. Complex dynamics in higher dimension. II. *Modern methods in complex analysis* (Princeton, NJ, 1992), 135-182, *Ann. of Math. Stud.* **137**, Princeton Univ. Press, Princeton, NJ, 1995.

[12] V. Guedj. Dynamics of polynomial mappings on $\mathbb{C}^2$. *Amer. J. Math.* **124** (2002), 75-106.

[13] S-M. Heinemann. Julia sets for holomorphic endomorphisms of $\mathbb{C}^n$. *Ergodic Theory Dynam. Syst.* **16** (1996), 1275-1296.

[14] S-M. Heinemann. Julia sets of skew products in $\mathbb{C}^2$. *Kyushu J. Math.* **52** (1998), 299-329.

[15] M. Jonsson. Dynamics of polynomial skew products on $\mathbb{C}^2$. *Math. Ann.* **314** (1999), 403-447.

[16] K. Ueno. Symmetries of Julia sets of nondegenerate polynomial skew products on $\mathbb{C}^2$. *Michigan Math. J.* **59** (2010), 153-168.

[17] K. Ueno. Weighted Green functions of nondegenerate polynomial skew products on $\mathbb{C}^2$. *Discrete Contin. Dyn. Syst.* **31** (2011), 985-996.

[18] K. Ueno. Fiberwise Green functions of skew products semiconjugate to some polynomial products on $\mathbb{C}^2$. *Kodai Math. J.* **35** (2012), 344-356.

[19] K. Ueno. Symmetries of Julia sets of polynomial skew products on $\mathbb{C}^2$. Submitted.

Toba National College of Maritime Technology, Mie 517-8501, Japan
E-mail address: `kueno@toba-cmt.ac.jp`